

CORRECTION

Open Access



# Correction: Oncological risk of proximal gastrectomy for proximal advanced gastric cancer after neoadjuvant chemotherapy

Yonghe Chen<sup>1,2,3</sup>, Xiaojiang Chen<sup>4</sup>, Yi Lin<sup>1</sup>, Shenyang Zhang<sup>5</sup>, Zhiwei Zhou<sup>4\*</sup> and Junsheng Peng<sup>1,2,3\*</sup>

**Correction:** *BMC Cancer* 24, 255 (2024)  
<https://doi.org/10.1186/s12885-024-11993-5>

Following publication of the original article [1], the authors reported an error in Fig. 1, specifically in colour spectrum within the heatmap located in the figure's lower right quadrant. The incorrect and the correct figures are supplied in this correction article.

The original article can be found online at <https://doi.org/10.1186/s12885-024-11993-5>.

\*Correspondence:

Zhiwei Zhou

[zhouzhw@sysucc.org.cn](mailto:zhouzhw@sysucc.org.cn)

Junsheng Peng

[pengjsh@mail.sysu.edu.cn](mailto:pengjsh@mail.sysu.edu.cn)

<sup>1</sup> Department of General Surgery, The Sixth Affiliated Hospital, Sun Yat-Sen University, 26 Yuancun Erheng Road, Guangzhou 510655, China

<sup>2</sup> Guangdong Provincial Key Laboratory of Colorectal and Pelvic Floor Diseases, The Sixth Affiliated Hospital, Sun Yat-Sen University, Guangzhou 510655, China



<sup>3</sup> Biomedical Innovation Center, The Sixth Affiliated Hospital, Sun Yat-Sen University, Guangzhou 510655, China

<sup>4</sup> Department of Gastric Surgery, Sun Yat-Sen University Cancer Center, Guangzhou 510060, China

<sup>5</sup> Department of Pathology, The Sixth Affiliated Hospital, Sun Yat-Sen University, Guangzhou 510655, China



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

**Incorrect Fig. 1****Corrected Fig. 1**

**Fig. 1** Q1 **(a)** This data map illustrates the metastasis rate of perigastric lymph nodes in proximal gastric cancer patients after neoadjuvant chemotherapy. Lymph nodes surrounding the proximal part of the stomach, such as No. 1/2/3/7, exhibit the highest metastasis rate (17.3% ~ 30%). Key distal lymph nodes, including No. 5/6/12a, have a collective metastasis rate of 10%. **(b)** This heatmap provides a visualization of the metastasis rate of grouped perigastric lymph nodes

The original article [1] has been corrected.

Published online: 06 March 2024

**Reference**

- Chen Y, Chen X, Lin Y, et al. Oncological risk of proximal gastrectomy for proximal advanced gastric cancer after neoadjuvant chemotherapy. BMC Cancer. 2024;24:255. <https://doi.org/10.1186/s12885-024-11993-5>.